

Experimental search for DCCs using Robust Observables

Debsankar Mukhopadhyay^{1 2 3}
for WA98 Collaboration
Variable Energy Cyclotron Centre,
1/AF, Bidhan Nagar, Calcutta - 700064,
India.

Abstract

We have analyzed the joint charged particle to photon density correlations in WA98 experiment at CERN-SPS in order to study the formation of the chirally disoriented vacuum in Pb + Pb collisions at 158 AGeV. The method of analysis is based on the event-by-event estimation of a set of robust observables which are the ratios of normalized bivariate factorial moments corrected for many efficiency factors. This is found to be a very powerful technique in identifying, within the limits of the detector resolution, the events with the localized charged particle to photon density fluctuations.

In high energy heavy ion collisions, hot fireballs may be produced with interiors having anomalous chiral order parameters. Eventually, these conjectured domains of disoriented chiral condensate(DCC) would decay by radiating pions with distinctive isospin distribution, following an inverse square-root law [1]

$$\left(\frac{dP}{df}\right)_{DCC} = \frac{1}{2\sqrt{f}}, \quad (1)$$

where f is the neutral pion fraction, given by

$$f = \frac{N_{\pi^0}}{N_{\pi^0} + N_{\pi^+} + N_{\pi^-}} \sim \frac{N_{\gamma}/2}{N_{\gamma}/2 + N_{ch}} \quad (2)$$

¹E-Mail: dsm@vecdec.veccal.ernet.in

²Financially supported by Council of Scientific and Industrial Research, Ministry of Human Resource Development Group, Government of India.

³Talk presented at the 3rd International Conference on Physics and Astrophysics of Quark-Gluon Plasma, March 17 - 21, Jaipur, India.

This is markedly different from the generic distribution, which is governed by a binomial distribution sharply peaked at $f \approx 1/3$.

Therefore, an experimental study of event by event charged to neutral density correlation would shed light on this exotic phenomenon.

WA98[2] is the only heavy ion experiment at CERN SPS which has initiated the ‘vacuum engineering’ in Pb + Pb collisions at 158 AGeV. The multiplicities of the charged particles are measured by a Silicon Pad Multiplicity Detector (SPMD) within a pseudorapidity coverage of $2.35 \leq \eta \leq 3.75$. Photon multiplicities are measured in the range of $2.4 \leq \eta \leq 4.4$ by a highly granular Photon Multiplicity Detector (PMD). The efficiency of detecting a charged particle in the active area of the detector is $\sim 99\%$, while the average photon counting efficiency is $70\% - 80\%$ depending on centrality, with a 35% contamination of showering hadrons. The centrality of the collision is determined by a Mid Rapidity Calorimeter (MIRAC) and a Zero Degree Calorimeter (ZDC) by respective measurement of the transverse and forward energies.

In addition to the detection efficiencies for charged particles and photon measurements, the following probability factors also exist because of finite acceptance of the detectors:

- (a) ϵ_{ch} : Probability of observing a $\pi^\pm$ within the detector acceptance, (b) ϵ_0 : Probability that none of the photons from a π^0 falls in the detector, (c) ϵ_1 : Probability that one of the γ ’s from a π^0 comes within the detector coverage, (d) ϵ_2 : Probability of observing both the γ ’s from a π^0 within the detector acceptance.

These ϵ factors along with the detection efficiencies and the background effects[3] put some serious challenges for us in this game.

Nevertheless, a sophisticated analysis technique, based on the method of the normalized factorial moments, has been developed [4] to estimate, within limits, the inclusive charged to neutral density correlations. In this formalism, the acceptance factors, ϵ_{ch} and ϵ_i ($i=0,1,2$), which can not be directly estimated from the data, are incorporated in the generating function of the bi-variate factorial moments. The resulting factorial moments contain the efficiency factors.

It can be shown[4] that the proper combinations of these factorial moments eliminate the acceptance factors and formulate the following ‘robust’ observables [4]:

$$r_{i,1} = \frac{F_{i,1}}{F_{i+1,0}}, \quad (3)$$

where F_i and $F_{i,j}$ are given by:

$$F_i \equiv \frac{\langle N(N-1) \dots (N-i+1) \rangle}{\langle N \rangle^i} \quad (4)$$

$$F_{i,j} = \frac{\langle n_{ch}(n_{ch}-1) \dots (n_{ch}-i+1) n_\gamma(n_\gamma-1) \dots (n_\gamma-j+1) \rangle}{\langle n_{ch} \rangle^i \langle n_\gamma \rangle^j}. \quad (5)$$

In the equations cited above, the angular brackets $\langle \dots \rangle$ denote the averages over a large number of events and N is the individual event multiplicity.

Moreover, one finds that for all $i \geq 1$

$$\begin{aligned} r_{i,1}(\text{generic}) &= 1, \\ r_{i,1}(\text{DCC}) &= \frac{1}{i+1} \end{aligned} \tag{6}$$

in the two cases.

The above formalism developed by the MiniMax Collaboration is not sensitive to the localized fluctuations in single events. In order to make it suitable to identify the single event fluctuation structures, we have to calculate the robust observables on an event-by-event basis. In that case the angular brackets in the numerators and denominators of equations (4) - (5) should denote the averages over a considerable number of bins in a single event phase space distributions of $N_\gamma - N_{ch}$. Now, the question comes whether one can use equation (6) to distinguish between a pure DCC and pure Generic event. The issue here is the size and correlation length of a DCC domain [5]. For example, if there is a single domain covering the entire detector acceptance, then one would expect that each bin (up to statistical fluctuations) would have the same orientation. On the other hand, if the DCC were produced throughout the acceptance but the correlation length is comparable to the bin size, then one might expect a $1/\sqrt{f}$ distribution (contaminated by whatever background is present) more or less bin by bin and in that case equation (6) can be used to distinguish a DCC and a generic event. The purpose of the multi-resolution analysis with moments or wavelets is to find out this type of correlations. Hence, in the WA98 environment we shall compute the robust observables event - by - event and try to explore their properties. This is important for using them as a distinctive tool for a DCC and generic event.

To analyze the data, we have selected the central events which correspond to about top 10% of the minimum bias cross-section. The selection of the central events confine us to the high detection efficiency regions specially for photons. The joint charged particle / photon distributions were constructed by using the overlap region $3.25 \leq \eta \leq 3.75$ of PMD and SPMD for the full azimuthal coverage. We have partitioned the whole azimuthal coverage into four different types of bins, namely, (a) 32 bins: $\Delta\phi = 11.25^\circ$, (b) 16 bins: $\Delta\phi = 22.50^\circ$, (c) 8 bins: $\Delta\phi = 45.00^\circ$, (d) 4 bins: $\Delta\phi = 90.00^\circ$, in order to achieve the multi-resolution binning of the available $\eta - \phi$ region.

In order to search for any anomalous charged to neutral particle density fluctuation, we have computed the robust observables for the following three sets of events: (1) generic events, (2) simulated exotic events and (3) experimental data.

The generic events are generated by the Monte-Carlo code VENUS [6]. The detector effects are introduced in VENUS in order to generate data-like generic events.

The simulation of the exotic events has been done by modifying the output of the VENUS event generator to include the fluctuations in the relative production of the charged particles and neutral pions. We assume that a single DCC domain

translates to the percent of VENUS pions that gets affected. The identities of these affected pions are altered by choosing a certain value of f randomly from the distribution given by (1). For the present purpose, we have simulated the exotic events in such a way that all of the pions within $3.0 \leq \eta \leq 4.0$ and $0^\circ \leq \phi \leq 45^\circ$ are supposed to come from a DCC domain. This is an example of a very large domain of DCC which may not be created in experiments but is very useful to test the sensitivity of the method towards the localized charged to neutral fluctuations.

The results of the analysis are summerized in Table -1 for the above three cases. The entire analysis has been done with a large number of events so that the statistical errors are small. In case of the data and simulated generic events, the distributions of $r_{i,1}$ are peaked around 1.00. But the mean value of the distributions shift to a lower value in case of the simulated exotic events. This indicates the sensitivity of the method to disentangle exotic events accompanied with a large domain of DCC from the generic samples. Moreover, the difference of the data and the simulated exotic events indicates the probability of the formation of smaller and/or multiple domains of DCC. Further simulations in this direction are in progress. The sensitiveness of the higher order moments (say, $r_{3,1}$) to the localized fluctuations is remarkable within the limits of the so called ‘empty bin effects’. The root mean square values of the distribution of $r_{i,1}$, for all resolutions, are higher for data than the simulated normal events. Also, the values of σ increases with decreasing bin size. This indicates that one should not go up to the smallest bin samples to avoid the trivial statistical effects.

Table - 1

The average values of $r_{i,1}$ for the simulated normal events, simulated exotic events and data.

Event type	No. of Bins	$r_{1,1} \pm \sigma$	$r_{2,1} \pm \sigma$	$r_{3,1} \pm \sigma$
Normal	4	1.001 ± 0.02	0.997 ± 0.02	1.008 ± 0.03
Exotic		0.845 ± 0.04	0.806 ± 0.04	0.742 ± 0.05
Data		1.009 ± 0.02	1.019 ± 0.03	1.031 ± 0.05
Normal	8	1.001 ± 0.01	1.004 ± 0.02	1.009 ± 0.04
Exotic		0.852 ± 0.04	0.848 ± 0.04	0.801 ± 0.05
Data		1.012 ± 0.03	1.027 ± 0.05	1.044 ± 0.07
Normal	16	1.002 ± 0.02	1.007 ± 0.04	1.010 ± 0.06
Exotic		0.889 ± 0.04	0.853 ± 0.04	0.831 ± 0.05
Data		1.021 ± 0.05	1.046 ± 0.07	1.080 ± 0.11
Normal	32	1.002 ± 0.04	1.009 ± 0.08	1.090 ± 0.12
Exotic		0.901 ± 0.04	0.889 ± 0.05	0.821 ± 0.08
Data		1.028 ± 0.05	1.067 ± 0.11	1.130 ± 0.19

A closer look into the data indicates a few events having the values of $r_{i,1} > 3\sigma$, where σ denotes the root mean square value of the concerned distribution. These events may be interesting with the indication of considerable charged particle

/ photon density fluctuations in the azimuthal space. But a concrete inference needs more studies on the detector effects as the data of both the PMD and SPMD are in a preliminary stage. This is, therefore, evident that in the first attempt of event-by-event computation of the robust observables, we have established the suitability of the moment analysis in the DCC search. It is found that the data contains more fluctuations than the simulation and for both the data and simulation σ increases with decreasing bin size. Also, the preliminary data indicates the presence of remarkable charged to neutral density fluctuations in some events. These results, thus, demand a more careful analysis of the detector effects for a confident conclusion.

References

- [1] J. D. Bjorken, K. L. Kowalski and C. C. Taylor, SLAC-PUB-6109, Proc. of Les Rencontres de la Vallée D' Aoste, La Thuile, 1993, ed. M. Greco, Editions Frontier, p. 507 (1993); K. Rajagopal and F. Wilczek, Nucl. Phys. B399 (1993)395.
- [2] Proposal for a Large Acceptance Hadron and Photon Spectrometer, CERN/SPSLC 91-17.
- [3] T. K. Nayak et al (WA98 Collaboration), these proceedings.
- [4] T. C. Brooks et al (MiniMax Collaboration), hep-ph/9609375.
- [5] C. C. Taylor, Private Communications.
- [6] K. Werner, Phys. Rev. Lett. 62(1989)2460.